

NDMI012: Combinatorics and Graph Theory 2


Lecture #10 Extremal combinatorics

Irena Penev

1 Turán's theorem

Given a positive integer n and a graph H , an n -vertex graph G without an H subgraph is *extremal* (for the property of not containing H as a subgraph) if it has the largest possible number of edges among all n -vertex graphs without an H subgraph; $\text{ex}(n, H)$ is the number of edges of an extremal n -vertex graph without an H subgraph. In other words, $\text{ex}(n, H)$ is the maximum number of edges that an n -vertex graph that does not contain H as a subgraph can have.

Obviously, any extremal graph G without an H subgraph is “edge-maximal” without an H subgraph, i.e. any graph obtained from G by adding one or more edges to it, contains H as a subgraph. The converse, however, does not hold in general: it is possible that a graph is edge-maximal without an H subgraph, without being extremal. For example, $2K_2$ is a four-vertex edge-maximal graph without a P_4 subgraph,¹ but it is not extremal: indeed, $K_{1,3}$ also has four vertices and no P_4 subgraph, and it has more edges than $2K_2$.




The following was proven in Combinatorics & Graphs Theory 1.

Mantel's theorem. *For any positive integer n , we have that $\text{ex}(n, K_3) = \lfloor \frac{n^2}{4} \rfloor$, and moreover, $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$ is an extremal n -vertex graph without a K_3 subgraph.*

¹As usual, P_4 is the four-vertex (and three-edge) path.

Mantel's theorem is a special case of Turán's theorem, to which we now turn.


For a positive integer r , a *complete r -partite graph* is a graph G whose vertex set can be partitioned into r (possibly empty) stable sets (called *parts*), pairwise complete to each other. For example, the graph below is complete 3-partite, with parts S_1, S_2, S_3 .



A *complete multipartite graph* is any graph that is complete r -partite for some r .

The *r -partite Turán graph on n vertices*, denoted by $T_r(n)$, is the complete r -partite graph on n vertices, in which the sizes of any two parts differ by at most one (so, each part is of size $\lfloor \frac{n}{r} \rfloor$ or $\lceil \frac{n}{r} \rceil$); $t_r(n)$ is the number of edges of $T_r(n)$. We note that the complete 3-partite graph above is in fact the graph $T_3(8)$.

Recall that *duplicating a vertex x* of a graph G produces a supergraph $G \circ x$ by adding to G a vertex x' and making it adjacent to all the neighbors of x in G , and to no other vertices of G (in particular, x and x' are nonadjacent in $G \circ x$). An example is shown below.



Obviously, $\omega(G \circ x) = \omega(G)$, i.e. G contains K_{r+1} is a subgraph if and only if $G \circ x$ does.

Turán's theorem. *Let n and r be positive integers. Then $ex(n, K_{r+1}) = t_r(n)$, and furthermore, $T_r(n)$ is the unique (up to isomorphism) extremal n -vertex graph without a K_{r+1} subgraph.*

Proof. We may assume that $r < n$, for otherwise, $T_r(n) \cong K_n$, and the result is immediate.

It is clear that $T_r(n)$ is an n -vertex graph without a K_{r+1} subgraph. Now, let G be any n -vertex extremal graph without a K_{r+1} subgraph. We must show that $G \cong T_r(n)$.

Claim. G is a complete multipartite graph.

Proof of the Claim. Suppose otherwise. Then non-adjacency is not an equivalence relation on $V(G)$, and it follows that there exist pairwise distinct vertices $y_1, x, y_2 \in V(G)$ such that $y_1x, xy_2 \notin E(G)$, but $y_1y_2 \in E(G)$. If $d_G(y_1) > d_G(x)$, then $G_1 := (G \setminus x) \circ y_1$ is an n -vertex graph that does not contain K_{r+1} as a subgraph, and $|E(G_1)| > |E(G)|$, contrary to the fact that G is extremal.

So, $d_G(y_1) \leq d_G(x)$, and similarly, $d_G(y_2) \leq d_G(x)$. Now, let G' be the graph obtained from $G \setminus \{y_1, y_2\}$ by duplicating x twice. Then G' is an n -vertex graph with no K_{r+1} subgraph, and (since $y_1y_2 \in E(G)$) we have that $|E(G')| = |E(G)| - (d_G(y_1) + d_G(y_2)) + 1 + 2d_G(x) \geq |E(G)| + 1$, contrary to the fact that G is extremal. This proves the Claim. ■

Now, using the Claim, we fix a partition $(S_1, \dots, S_k)$ of $V(G)$ into non-empty stable sets, pairwise complete to each other. Clearly, G contains K_k as a subgraph,² and so $k \leq r$. Suppose that $k < r$. Then since $r < n$, at least one of the sets $S_1, \dots, S_k$ has more than one vertex; by symmetry, we may assume that $|S_k| \geq 2$. Fix $a \in S_k$. Then consider the graph G' obtained from G by adding edges between a and all vertices of $S_k \setminus \{a\}$; then G' is a complete $(k+1)$ -partite graph, it does not contain K_{r+1} as a subgraph (because $k < r$), and it has more edges than G , contrary to the fact that G is extremal. So, $k = r$.

It remains to show that any two of $S_1, \dots, S_r$ differ in size by at most one (this will imply that $G \cong T_r(n)$). Suppose otherwise. By symmetry, we may assume that $|S_1| \geq |S_2| + 2$. Now, fix a vertex $a \in S_1$, and let G' be the graph obtained by first deleting all edges between a and S_2 , and then adding all edges between a and $S_1 \setminus \{a\}$. (This effectively “moves” a into S_2 .) Now G' is still a complete r -partite graph on n vertices, and it does not contain K_{r+1} as a subgraph. Furthermore, since $|S_1| \geq |S_2| + 2$, we see that $|E(G')| \geq |E(G)| + 1$. But this contradicts the fact that G is extremal. □

2 The Erdős-Ko-Rado theorem

Suppose we are given positive integers r and n , and we want to select a maximum number of pairwise intersecting r -element subsets of $\{1, \dots, n\}$. What is this maximum number? For $r > \frac{n}{2}$, any two r -element subsets of $\{1, \dots, n\}$ intersect, and there are $\binom{n}{r}$ such subsets. How about if $r \leq \frac{n}{2}$? In that case, we can consider all r -element subsets of $\{1, \dots, n\}$ that contain n ; there are $\binom{n-1}{r-1}$ such subsets, and obviously, they pairwise intersect. As the following theorem shows, this is in fact best possible.

²Indeed, we just take one vertex from each S_i , and we obtain a clique of size k .

The Erdős-Ko-Rado theorem. *Let r and n be positive integers such that $r \leq \frac{n}{2}$. Then there are at most $\binom{n-1}{r-1}$ pairwise intersecting r -element subsets of $\{1, \dots, n\}$.*

Proof. Let $A_1, \dots, A_m$ be pairwise distinct and pairwise intersecting r -element subsets of $\{1, \dots, n\}$. We must show that $m \leq \binom{n-1}{r-1}$.

Let c be the number of ordered pairs (C, A) , where

- C is a directed cycle with vertex set $\{1, \dots, n\}$;³
- A is an r -vertex directed subpath of C ;
- $V(A) = A_i$ for some $i \in \{1, \dots, m\}$.

Now we count in two ways, as follows. On the one hand, we can form an ordered pair (C, A) by first selecting one of the sets $A_1, \dots, A_m$ (we have m choices), then ordering its vertices to form a directed path (there are $r!$ choices), and then ordering the remaining $n - r$ vertices to complete the cycle C (there are $(n - r)!$ choices). So,

$$c = mr!(n - r)!.$$

We now count in another way. First, there are $(n - 1)!$ ways of ordering $\{1, \dots, n\}$ to obtain a directed cycle C . Next, we claim that for fixed C , there are at most r directed subpaths of C that correspond to one of A_i 's. Indeed, suppose the subpath $a_1, a_2, \dots, a_r$ of C corresponds to one of $A_1, \dots, A_m$. Then for any other subpath of C corresponding to one of $A_1, \dots, A_m$ (and therefore containing at least one of $a_1, \dots, a_r$), there exists some $i \in \{1, \dots, n - 1\}$ such that either a_i is the terminal vertex of the path, or a_{i+1} is the initial vertex of the path; but since $r \leq \frac{n}{2}$, the r -vertex subpath terminating at a_i and the r -vertex subpath starting at a_{i+1} have no vertices in common, and so at most one of them can correspond to one of $A_1, \dots, A_m$. Thus, in addition to $a_1, \dots, a_r$, there are at most $r - 1$ subpaths of C corresponding to one of $A_1, \dots, A_m$; in total, at most r subpaths of C correspond to one of $A_1, \dots, A_m$. This proves that

$$c \leq (n - 1)!r.$$

We now have that

$$mr!(n - r)! = c \leq (n - 1)!r,$$

and so

$$m \leq \frac{(n-1)!r}{r!(n-r)!} = \binom{n-1}{r-1},$$

which is what we needed to show. □

³Vertices $1, \dots, n$ need **not** appear in that order on the cycle.

3 The Sunflower lemma

A *sunflower* is a family (i.e. collection) $\mathcal{S}$ of sets (called *petals*) such that there exists a set S (called a *kernel*) with the property that for all distinct $S_1, S_2 \in \mathcal{S}$, we have that $S_1 \cap S_2 = S$.

The Sunflower lemma (Erdős-Rado). *Let ℓ and p be positive integers, and let $\mathcal{A}$ be a family of sets such that*

- $|A| \leq \ell$ for all $A \in \mathcal{A}$, and
- $|\mathcal{A}| > (p-1)^\ell \ell!$.

Then there exists a sunflower $\mathcal{S} \subseteq \mathcal{A}$ with p petals.

Proof. We assume inductively that the lemma is true for smaller values of ℓ . More precisely, we assume that for all positive integers $\ell' < \ell$, and all families $\mathcal{A}'$ of sets such that

- $|A| \leq \ell'$ for all $A \in \mathcal{A}'$, and
- $|\mathcal{A}'| > (p-1)^{\ell'} \ell'!$,

there exists a sunflower $\mathcal{S}' \subseteq \mathcal{A}'$ with p petals.

Note that $|\mathcal{A}| \geq p$; so, if $p \leq 2$, then any p elements of $\mathcal{A}$ form a sunflower with p petals, and we are done. So, we may assume that $p \geq 3$. Next, suppose that $\ell = 1$. Then $|A| \leq 1$ for all $A \in \mathcal{A}$ and $|\mathcal{A}| > p-1$. We then take any p elements of $\mathcal{A}$, and we observe that they form a sunflower (with an empty kernel). So, from now on, we assume that $\ell \geq 2$.

Let $\mathcal{D} \subseteq \mathcal{A}$ be a collection of pairwise disjoint sets, with $|\mathcal{D}|$ chosen maximum. If $|\mathcal{D}| \geq p$, then any p elements of $\mathcal{D}$ form a sunflower (with an empty kernel), and we are done. So assume that $|\mathcal{D}| < p$. Let $D = \bigcup \mathcal{D}$; then $|D| \leq |\mathcal{D}| \ell \leq (p-1)\ell$. Furthermore, since $|\mathcal{A}| \geq \ell \geq 2$, the maximality of $\mathcal{D}$ guarantees that $\mathcal{D}$ contains at least one non-empty set, and so $D \neq \emptyset$.

Claim. There exists some $d \in D$ such that d belongs to more than $(p-1)^{\ell-1}(\ell-1)!$ elements of $\mathcal{A}$.

Proof of the Claim. We consider two cases: when $\emptyset \in \mathcal{A}$, and when this is not the case.

Suppose first that $\emptyset \notin \mathcal{A}$ (and consequently, $\emptyset \notin \mathcal{D}$). Then every element of $\mathcal{A}$ intersects D : indeed, since $\emptyset \notin \mathcal{D}$, we know that every element of $\mathcal{D}$ intersects D , and by the maximality of $\mathcal{D}$, every element of $\mathcal{A} \setminus \mathcal{D}$ intersects D . But then by the Pigeonhole Principle, some element of D belongs to at least

$$\left\lceil \frac{|\mathcal{A}|}{|D|} \right\rceil > \frac{(p-1)^\ell \ell!}{(p-1)\ell} = (p-1)^{\ell-1}(\ell-1)!,$$

elements of $\mathcal{A}$, which is what we needed.

Suppose now that $\emptyset \in \mathcal{A}$. Then the maximality of $\mathcal{D}$ guarantees that $\emptyset \in D$. Since $D \neq \emptyset$, it follows that $|\mathcal{D}| \geq 2$. Since $\emptyset \in \mathcal{D}$, we see that $|D| \leq (|\mathcal{D}| - 1)\ell \leq (p - 2)\ell$. Now by the maximality of $\mathcal{D}$, every element of $\mathcal{A} \setminus \{\emptyset\}$ intersects D . But then by the Pigeonhole Principle, some element of D belongs to at least

$$\begin{aligned} \left\lceil \frac{|\mathcal{A} \setminus \{\emptyset\}|}{|D|} \right\rceil &\geq \frac{(p-1)^\ell \ell!}{(p-2)^\ell} \\ &= (p-1)^{\ell-1}(\ell-1)! \frac{p-1}{p-2} \\ &> (p-1)^{\ell-1}(\ell-1)! \end{aligned}$$

elements of $\mathcal{A}$, which is what we needed. This proves the Claim. ■

Let $d \in D$ be as in the Claim, and set $\mathcal{A}' := \{A \setminus \{d\} \mid A \in \mathcal{A}, d \in A\}$. Then $|\mathcal{A}'| > (p-1)^{\ell-1}(\ell-1)!$; furthermore, $|A| \leq \ell-1$ for all $A \in \mathcal{A}'$. So, by the induction hypothesis, there exists a sunflower $\mathcal{S}' \subseteq \mathcal{A}'$ with p petals. Now, set $\mathcal{S} := \{A \cup \{d\} \mid A \in \mathcal{S}'\}$; then $\mathcal{S} \subseteq \mathcal{A}$ is a sunflower with p petals, and we are done. $\square$