## NDMI012: Combinatorics and Graph Theory 2

Lecture #10

# Extremal combinatorics

Irena Penev

May 5, 2021

• Turán's theorem;

- Turán's theorem;
- the Erdős-Ko-Rado theorem;

- Turán's theorem;
- the Erdős-Ko-Rado theorem;
- the Sunflower lemma.

Part I: Turán's theorem

## Part I: Turán's theorem

## Definition

Given a positive integer n and a graph H, an n-vertex graph G without an H subgraph is *extremal* (for the property of not containing H as a subgraph) if it has the largest possible number of edges among all n-vertex graphs without an H subgraph; ex(n, H) is the number of edges of an extremal n-vertex graph without an H subgraph.

• So, ex(n, H) is the maximum number of edges that an *n*-vertex graph that does not contain H as a subgraph can have. • Any extremal graph G without an H subgraph is "edge-maximal" without an H subgraph, i.e. any graph obtained from G by adding one or more edges to it, contains H as a subgraph.

- Any extremal graph G without an H subgraph is "edge-maximal" without an H subgraph, i.e. any graph obtained from G by adding one or more edges to it, contains H as a subgraph.
- The converse, however, does not hold in general: it is possible that a graph is edge-maximal without an *H* subgraph, without being extremal.

- Any extremal graph G without an H subgraph is "edge-maximal" without an H subgraph, i.e. any graph obtained from G by adding one or more edges to it, contains H as a subgraph.
- The converse, however, does not hold in general: it is possible that a graph is edge-maximal without an *H* subgraph, without being extremal.
  - For example,  $2K_2$  is a four-vertex edge-maximal graph without a  $P_4$  subgraph, but it is not extremal: indeed,  $K_{1,3}$  also has four vertices and no  $P_4$  subgraph, and it has more edges than  $2K_2$ .



#### Mantel's theorem

For any positive integer *n*, we have that  $ex(n, K_3) = \lfloor \frac{n^2}{4} \rfloor$ , and moreover,  $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$  is an extremal *n*-vertex graph without a  $K_3$  subgraph.

Proof. Combinatorics & Graphs 1.

#### Mantel's theorem

For any positive integer *n*, we have that  $ex(n, K_3) = \lfloor \frac{n^2}{4} \rfloor$ , and moreover,  $K_{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil}$  is an extremal *n*-vertex graph without a  $K_3$  subgraph.

Proof. Combinatorics & Graphs 1.

 Mantel's theorem is a special case of Turán's theorem, to which we now turn.

## Definition

For a positive integer r, a *complete r-partite graph* is a graph G whose vertex set can be partitioned into r (possibly empty) stable sets (called *parts*), pairwise complete to each other.



#### Definition

A complete multipartite graph is any graph that is complete r-partite for some r.

#### Definition

The *r*-partite Turán graph on *n* vertices, denoted by  $T_r(n)$ , is the complete *r*-partite graph on *n* vertices, in which the sizes of any two parts differ by at most one (so, each part is of size  $\lfloor \frac{n}{r} \rfloor$  or  $\lceil \frac{n}{r} \rceil$ );  $t_r(n)$  is the number of edges of  $T_r(n)$ .



Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.



• Duplicating a vertex x of a graph G produces a supergraph  $G \circ x$  by adding to G a vertex x' and making it adjacent to all the neighbors of x in G, and to no other vertices of G (in particular, x and x' are nonadjacent in  $G \circ x$ ).



Duplicating a vertex x of a graph G produces a supergraph G ∘ x by adding to G a vertex x' and making it adjacent to all the neighbors of x in G, and to no other vertices of G (in particular, x and x' are nonadjacent in G ∘ x).



Obviously, ω(G ∘ x) = ω(G), i.e. G contains K<sub>r+1</sub> is a subgraph iff G ∘ x does.



Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

Proof.



Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof.* We may assume that r < n, for otherwise,  $T_r(n) \cong K_n$ , and the result is immediate.



Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof.* We may assume that r < n, for otherwise,  $T_r(n) \cong K_n$ , and the result is immediate.

It is clear that  $T_r(n)$  is an *n*-vertex graph without a  $K_{r+1}$  subgraph.



Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof.* We may assume that r < n, for otherwise,  $T_r(n) \cong K_n$ , and the result is immediate.

It is clear that  $T_r(n)$  is an *n*-vertex graph without a  $K_{r+1}$  subgraph. Now, let G be any *n*-vertex extremal graph without a  $K_{r+1}$  subgraph. We must show that  $G \cong T_r(n)$ .

Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

Proof of the Claim.

Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

Proof of the Claim. Suppose otherwise.

Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

*Proof of the Claim.* Suppose otherwise. Then non-adjacency is not an equivalence relation on V(G),

Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

*Proof of the Claim.* Suppose otherwise. Then non-adjacency is not an equivalence relation on V(G), and it follows that there exist pairwise distinct vertices  $y_1, x, y_2 \in V(G)$  s.t.  $y_1x, xy_2 \notin E(G)$ , but  $y_1y_2 \in E(G)$ .

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

Proof of the Claim. Suppose otherwise. Then non-adjacency is not an equivalence relation on V(G), and it follows that there exist pairwise distinct vertices  $y_1, x, y_2 \in V(G)$  s.t.  $y_1x, xy_2 \notin E(G)$ , but  $y_1y_2 \in E(G)$ . If  $d_G(y_1) > d_G(x)$ , then  $G_1 := (G \setminus x) \circ y_1$  is an *n*-vertex graph that does not contain  $K_{r+1}$  as a subgraph, and  $|E(G_1)| > |E(G)|$ , contrary to the fact that G is extremal.

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

**Claim.** G is a complete multipartite graph.

Proof of the Claim. Suppose otherwise. Then non-adjacency is not an equivalence relation on V(G), and it follows that there exist pairwise distinct vertices  $y_1, x, y_2 \in V(G)$  s.t.  $y_1x, xy_2 \notin E(G)$ , but  $y_1y_2 \in E(G)$ . If  $d_G(y_1) > d_G(x)$ , then  $G_1 := (G \setminus x) \circ y_1$  is an *n*-vertex graph that does not contain  $K_{r+1}$  as a subgraph, and  $|E(G_1)| > |E(G)|$ , contrary to the fact that G is extremal. So,  $d_G(y_1) \leq d_G(x)$ , and similarly,  $d_G(y_2) \leq d_G(x)$ .

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Claim. G is a complete multipartite graph.

Proof of the Claim (continued). Reminder:  $y_1x, xy_2 \notin E(G)$  and  $y_1y_2 \in E(G)$ ;  $d_G(y_1), d_G(y_2) \leq d_G(x)$ .

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Claim. G is a complete multipartite graph.

Proof of the Claim (continued). Reminder:  $y_1x, xy_2 \notin E(G)$  and  $y_1y_2 \in E(G)$ ;  $d_G(y_1), d_G(y_2) \leq d_G(x)$ . Now, let G' be the graph obtained from  $G \setminus \{y_1, y_2\}$  by duplicating x twice.

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Claim. G is a complete multipartite graph.

Proof of the Claim (continued). Reminder:  $y_1x, xy_2 \notin E(G)$  and  $y_1y_2 \in E(G)$ ;  $d_G(y_1), d_G(y_2) \leq d_G(x)$ . Now, let G' be the graph obtained from  $G \setminus \{y_1, y_2\}$  by duplicating x twice. Then G' is an *n*-vertex graph with no  $K_{r+1}$  subgraph, and (since  $y_1y_2 \in E(G)$ ) we have that

 $|E(G')| = |E(G)| - (d_G(y_1) + d_G(y_2)) + 1 + 2d_G(x) \ge |E(G)| + 1$ , contrary to the fact that G is extremal. This proves the Claim.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Now, using the Claim, we fix a partition  $(S_1, \ldots, S_k)$  of V(G) into non-empty stable sets, pairwise complete to each other.

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Now, using the Claim, we fix a partition  $(S_1, \ldots, S_k)$  of V(G) into non-empty stable sets, pairwise complete to each other. Clearly, G contains  $K_k$  is a subgraph, and so  $k \leq r$ .

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Now, using the Claim, we fix a partition  $(S_1, \ldots, S_k)$  of V(G) into non-empty stable sets, pairwise complete to each other. Clearly, G contains  $K_k$  is a subgraph, and so  $k \leq r$ . Suppose that k < r.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Now, using the Claim, we fix a partition  $(S_1, \ldots, S_k)$  of V(G) into non-empty stable sets, pairwise complete to each other. Clearly, Gcontains  $K_k$  is a subgraph, and so  $k \leq r$ . Suppose that k < r. Then since r < n, at least one of the sets  $S_1, \ldots, S_k$  has more than one vertex; by symmetry, we may assume that  $|S_k| \geq 2$ . Fix  $a \in S_k$ .

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph.

Now, using the Claim, we fix a partition  $(S_1, \ldots, S_k)$  of V(G) into non-empty stable sets, pairwise complete to each other. Clearly, Gcontains  $K_k$  is a subgraph, and so  $k \le r$ . Suppose that k < r. Then since r < n, at least one of the sets  $S_1, \ldots, S_k$  has more than one vertex; by symmetry, we may assume that  $|S_k| \ge 2$ . Fix  $a \in S_k$ . Then we get a contradiction to the fact that G is extremal as follows (next slide; formal details: Lecture Notes).

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

Proof (continued).


Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

Proof (continued).



So, r = k.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ .

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one. Suppose otherwise.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one. Suppose otherwise. By symmetry, we may assume that  $|S_1| \ge |S_2| + 2$ .

Let *n* and *r* be positive integers. Then  $e_X(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one. Suppose otherwise. By symmetry, we may assume that  $|S_1| \ge |S_2| + 2$ . Now, fix a vetex  $a \in S_1$ , and "move" *a* from  $S_1$  to  $S_2$  (i.e. delete edges between *a* and  $S_2$ , and add edges between *a* and  $S_1 \setminus \{a\}$ ).

Let *n* and *r* be positive integers. Then  $e_x(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; *G* is *n*-vertex extremal graph without a  $K_{r+1}$  subgraph; *G* is a complete *r*-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one. Suppose otherwise. By symmetry, we may assume that  $|S_1| \ge |S_2| + 2$ . Now, fix a vetex  $a \in S_1$ , and "move" *a* from  $S_1$  to  $S_2$  (i.e. delete edges between *a* and  $S_2$ , and add edges between *a* and  $S_1 \setminus \{a\}$ ). This increases the number of edges without creating a  $K_{r+1}$  subgraph, contrary to the fact that *G* is extremal.

Let *n* and *r* be positive integers. Then  $ex(n, K_{r+1}) = t_r(n)$ , and furthermore,  $T_r(n)$  is the unique (up to isomorphism) extremal *n*-vertex graph without a  $K_{r+1}$  subgraph.

*Proof (continued).* Reminder: r < n; G is n-vertex extremal graph without a  $K_{r+1}$  subgraph; G is a complete r-partite graph with (non-empty) parts  $S_1, \ldots, S_r$ . WTS  $G \cong T_r(n)$ . It remains to show that any two of  $S_1, \ldots, S_r$  differ in size by at most one. Suppose otherwise. By symmetry, we may assume that  $|S_1| \ge |S_2| + 2$ . Now, fix a vetex  $a \in S_1$ , and "move" a from  $S_1$  to  $S_2$  (i.e. delete edges between a and  $S_2$ , and add edges between a and  $S_1 \setminus \{a\}$ ). This increases the number of edges without creating a  $K_{r+1}$  subgraph, contrary to the fact that G is extremal. So,  $G \cong T_r(n)$ .

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?
- For r > <sup>n</sup>/<sub>2</sub>, any two r-element subsets of {1,..., n} intersect, and there are (<sup>n</sup>/<sub>r</sub>) such subsets.

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?
- For r > n/2, any two r-element subsets of {1,..., n} intersect, and there are (n/r) such subsets.
- How about if  $r \leq \frac{n}{2}$ ?

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?
- For r > n/2, any two r-element subsets of {1,..., n} intersect, and there are (n/r) such subsets.
- How about if  $r \leq \frac{n}{2}$ ?
  - In that case, we can consider all *r*-element subsets of  $\{1, \ldots, n\}$  that contain *n*.
  - There are  $\binom{n-1}{r-1}$  such subsets, and obviously, they pairwise intersect.

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?
- For r > n/2, any two r-element subsets of {1,..., n} intersect, and there are (n/r) such subsets.
- How about if  $r \leq \frac{n}{2}$ ?
  - In that case, we can consider all *r*-element subsets of  $\{1, \ldots, n\}$  that contain *n*.
  - There are  $\binom{n-1}{r-1}$  such subsets, and obviously, they pairwise intersect.
  - As the following theorem shows, this is in fact best possible.

- Suppose we are given positive integers *r* and *n*, and we want to select a maximum number of pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .
  - What this this maximum number?
- For r > <sup>n</sup>/<sub>2</sub>, any two r-element subsets of {1,..., n} intersect, and there are (<sup>n</sup>/<sub>r</sub>) such subsets.
- How about if  $r \leq \frac{n}{2}$ ?
  - In that case, we can consider all *r*-element subsets of  $\{1, \ldots, n\}$  that contain *n*.
  - There are  $\binom{n-1}{r-1}$  such subsets, and obviously, they pairwise intersect.
  - As the following theorem shows, this is in fact best possible.

### The Erdős-Ko-Rado theorem

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

Let r and n be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting r-element subsets of  $\{1, \ldots, n\}$ .

Proof.

Let r and n be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting r-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (C, A), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Let r and n be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting r-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (*C*, *A*), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Now we count in two ways, as follows.

Let r and n be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting r-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (C, A), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Now we count in two ways, as follows. On the one hand, we can form an ordered pair (C, A) by first selecting one of the sets  $A_1, \ldots, A_m$  (we have *m* choices),

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (C, A), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Now we count in two ways, as follows. On the one hand, we can form an ordered pair (C, A) by first selecting one of the sets  $A_1, \ldots, A_m$  (we have *m* choices), then ordering its vertices to form a directed path (there are r! choices),

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (C, A), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Now we count in two ways, as follows. On the one hand, we can form an ordered pair (C, A) by first selecting one of the sets  $A_1, \ldots, A_m$  (we have *m* choices), then ordering its vertices to form a directed path (there are r! choices), and then ordering the remaining n - r vertices to complete the cycle *C* (there are (n - r)! choices).

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

*Proof.* Let  $A_1, \ldots, A_m$  be pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ . WTS  $m \leq \binom{n-1}{r-1}$ . Let *c* be the number of ordered pairs (C, A), where

- C is a directed cycle with vertex set {1,...,n};
  - vertics 1,..., n need **not** appear in that order on the cycle;
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ .

Now we count in two ways, as follows. On the one hand, we can form an ordered pair (C, A) by first selecting one of the sets  $A_1, \ldots, A_m$  (we have *m* choices), then ordering its vertices to form a directed path (there are *r*! choices), and then ordering the remaining n - r vertices to complete the cycle *C* (there are (n - r)! choices). So, c = mr!(n - r)!.

- C is a directed cycle with vertex set {1,..., n};
- A is an r-vertex directed subpath of C;

• 
$$V(A) = A_i$$
 for some  $i \in \{1, ..., m\}$ ;

$$c = mr!(n-r)!.$$

- *C* is a directed cycle with vertex set {1,..., *n*};
- A is an r-vertex directed subpath of C;
- $V(A) = A_i$  for some  $i \in \{1, \ldots, m\}$ ;
- c = mr!(n-r)!.

We now count in another way.

- *C* is a directed cycle with vertex set {1,..., *n*};
- A is an r-vertex directed subpath of C;

• 
$$V(A) = A_i$$
 for some  $i \in \{1, ..., m\};$ 

$$c = mr!(n-r)!.$$

We now count in another way. First, there are (n-1)! ways of ordering  $\{1, \ldots, n\}$  to obtain a directed cycle C.

- *C* is a directed cycle with vertex set {1,..., *n*};
- A is an r-vertex directed subpath of C;

• 
$$V(A) = A_i$$
 for some  $i \in \{1, ..., m\};$ 

$$c = mr!(n-r)!.$$

We now count in another way. First, there are (n-1)! ways of ordering  $\{1, \ldots, n\}$  to obtain a directed cycle *C*. WTS that for fixed *C*, there are at most *r* directed subpaths of *C* that correspond to one of  $A_i$ 's.

- *C* is a directed cycle with vertex set {1,..., *n*};
- A is an r-vertex directed subpath of C;

• 
$$V(A) = A_i$$
 for some  $i \in \{1, ..., m\}$ ;

$$c = mr!(n-r)!.$$

We now count in another way. First, there are (n-1)! ways of ordering  $\{1, \ldots, n\}$  to obtain a directed cycle *C*. WTS that for fixed *C*, there are at most *r* directed subpaths of *C* that correspond to one of  $A_i$ 's. Indeed, suppose the subpath  $a_1, a_2, \ldots, a_r$  of *C* corresponds to one of  $A_1, \ldots, A_m$ .





Then for any other subpath of *C* corresponding to one of  $A_1, \ldots, A_m$  (and therefore containing at least one of  $a_1, \ldots, a_r$ ), there exists some  $i \in \{1, \ldots, n-1\}$  s.t. either  $a_i$  is the terminal vertex of the path, or  $a_{i+1}$  is the initial vertex of the path;



Then for any other subpath of *C* corresponding to one of  $A_1, \ldots, A_m$  (and therefore containing at least one of  $a_1, \ldots, a_r$ ), there exists some  $i \in \{1, \ldots, n-1\}$  s.t. either  $a_i$  is the terminal vertex of the path, or  $a_{i+1}$  is the initial vertex of the path; but since  $r \leq \frac{n}{2}$ , at most one of those choices is possible.



Then for any other subpath of *C* corresponding to one of  $A_1, \ldots, A_m$  (and therefore containing at least one of  $a_1, \ldots, a_r$ ), there exists some  $i \in \{1, \ldots, n-1\}$  s.t. either  $a_i$  is the terminal vertex of the path, or  $a_{i+1}$  is the initial vertex of the path; but since  $r \leq \frac{n}{2}$ , at most one of those choices is possible. This proves that

$$c \leq (n-1)!r.$$

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

*Proof (continued).* Reminder:  $A_1, \ldots, A_m$  are pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ ;  $mr!(n-r)! = c \leq (n-1)!r$ .

Let *r* and *n* be positive integers s.t.  $r \leq \frac{n}{2}$ . Then there are at most  $\binom{n-1}{r-1}$  pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ .

*Proof (continued).* Reminder:  $A_1, \ldots, A_m$  are pairwise distinct and pairwise intersecting *r*-element subsets of  $\{1, \ldots, n\}$ ;  $mr!(n-r)! = c \le (n-1)!r$ . So, (n-1)!r.

$$m \leq \frac{(n-1)!r}{r!(n-r)!} = \binom{n-1}{r-1},$$

which is what we needed to show.

Part III: The Sunflower lemma

## Part III: The Sunflower lemma

#### Definition

A sunflower is a family (i.e. collection)  $\mathscr{S}$  of sets (called *petals*) s.t. there exists a set S (called a *kernel*) with the property that for all distinct  $S_1, S_2 \in \mathscr{S}$ , we have that  $S_1 \cap S_2 = S$ .

• A sunflower  $\mathscr{S} = \{S_1, \ldots, S_k\}$  with kernel S:



# Definition

A sunflower is a family (i.e. collection)  $\mathscr{S}$  of sets (called *petals*) s.t. there exists a set S (called a *kernel*) with the property that for all distinct  $S_1, S_2 \in \mathscr{S}$ , we have that  $S_1 \cap S_2 = S$ .

#### Definition

A sunflower is a family (i.e. collection)  $\mathscr{S}$  of sets (called *petals*) s.t. there exists a set S (called a *kernel*) with the property that for all distinct  $S_1, S_2 \in \mathscr{S}$ , we have that  $S_1 \cap S_2 = S$ .

# The Sunflower lemma [Erdős-Rado]

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

Proof.
#### Definition

A sunflower is a family (i.e. collection)  $\mathscr{S}$  of sets (called *petals*) s.t. there exists a set S (called a *kernel*) with the property that for all distinct  $S_1, S_2 \in \mathscr{S}$ , we have that  $S_1 \cap S_2 = S$ .

# The Sunflower lemma [Erdős-Rado]

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof.* We assume inductively that the lemma is true for smaller values of  $\ell$ .

#### Definition

A sunflower is a family (i.e. collection)  $\mathscr{S}$  of sets (called *petals*) s.t. there exists a set S (called a *kernel*) with the property that for all distinct  $S_1, S_2 \in \mathscr{S}$ , we have that  $S_1 \cap S_2 = S$ .

## The Sunflower lemma [Erdős-Rado]

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof.* We assume inductively that the lemma is true for smaller values of  $\ell$ . If  $p \leq 2$  or  $\ell = 1$ , then it's easy (details: Lecture Notes). So, we assume that  $p \geq 3$  and  $\ell \geq 2$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\mathscr{D}\subseteq\mathscr{A}$  be a collection of pairwise disjoint sets, with  $|\mathscr{D}|$  chosen maximum.

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets, with  $|\mathscr{D}|$  chosen maximum. If  $|\mathscr{D}| \ge p$ , then any p elements of  $\mathscr{D}$  form a sunflower (with an empty kernel), and we are done.

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets, with  $|\mathscr{D}|$  chosen maximum. If  $|\mathscr{D}| \ge p$ , then any p elements of  $\mathscr{D}$  form a sunflower (with an empty kernel), and we are done. So assume that  $|\mathscr{D}| < p$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets, with  $|\mathscr{D}|$  chosen maximum. If  $|\mathscr{D}| \ge p$ , then any p elements of  $\mathscr{D}$  form a sunflower (with an empty kernel), and we are done. So assume that  $|\mathscr{D}| < p$ . Let  $D = \bigcup \mathscr{D}$ ; then  $|D| \le |\mathscr{D}|\ell \le (p-1)\ell$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ .

Let  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets, with  $|\mathscr{D}|$  chosen maximum. If  $|\mathscr{D}| \ge p$ , then any p elements of  $\mathscr{D}$  form a sunflower (with an empty kernel), and we are done. So assume that  $|\mathscr{D}| < p$ . Let  $D = \bigcup \mathscr{D}$ ; then  $|D| \le |\mathscr{D}|\ell \le (p-1)\ell$ . Furthermore, since  $|\mathscr{A}| \ge \ell \ge 2$ , the maximality of  $\mathscr{D}$  guarantees that  $\mathscr{D}$  contains at least one non-empty set, and so  $D \ne \emptyset$ .

Proof of the Claim.

*Proof of the Claim.* We consider the case when  $\emptyset \notin \mathscr{A}$  (see the Lecture Notes for the other case).

*Proof of the Claim.* We consider the case when  $\emptyset \notin \mathscr{A}$  (see the Lecture Notes for the other case).

Then every element of  $\mathscr{A}$  intersects D: indeed, since  $\emptyset \notin \mathscr{D}$ , we know that every element of  $\mathscr{D}$  intersects D, and by the maximality of  $\mathscr{D}$ , every element of  $\mathscr{A} \setminus \mathscr{D}$  intersects D.

*Proof of the Claim.* We consider the case when  $\emptyset \notin \mathscr{A}$  (see the Lecture Notes for the other case).

Then every element of  $\mathscr{A}$  intersects D: indeed, since  $\emptyset \notin \mathscr{D}$ , we know that every element of  $\mathscr{D}$  intersects D, and by the maximality of  $\mathscr{D}$ , every element of  $\mathscr{A} \setminus \mathscr{D}$  intersects D. But then by the Pigeonhole Principle, some element of D belongs to at least

$$\left\lceil rac{|\mathscr{A}|}{|D|} 
ight
ceil > rac{(p-1)^\ell \ell!}{(p-1)\ell} = (p-1)^{\ell-1} (\ell-1)!,$$

elements of  $\mathcal{A}$ , which is what we needed. This proves the Claim.

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $d \in D$  be as in the Claim,

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $d \in D$  be as in the Claim, and set  $\mathscr{A}' := \{A \setminus \{d\} \mid A \in \mathscr{A}, d \in A\}.$ 

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $d \in D$  be as in the Claim, and set  $\mathscr{A}' := \{A \setminus \{d\} \mid A \in \mathscr{A}, d \in A\}$ . Then  $|\mathscr{A}'| > (p-1)^{\ell-1}(\ell-1)!$ , and  $|A| \leq \ell - 1 \ \forall A \in \mathscr{A}'$ .

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $d \in D$  be as in the Claim, and set  $\mathscr{A}' := \{A \setminus \{d\} \mid A \in \mathscr{A}, d \in A\}$ . Then  $|\mathscr{A}'| > (p-1)^{\ell-1}(\ell-1)!$ , and  $|A| \leq \ell - 1 \ \forall A \in \mathscr{A}'$ . So, by the ind. hyp.,  $\exists$  sunflower  $\mathscr{S}' \subseteq \mathscr{A}'$  with p petals.

Let  $\ell$  and p be positive integers, and let  $\mathscr{A}$  be a family of sets s.t.

- $|A| \leq \ell$  for all  $A \in \mathscr{A}$ , and
- $|\mathscr{A}| > (p-1)^{\ell} \ell!$ .

Then there exists a sunflower  $\mathscr{S} \subseteq \mathscr{A}$  with p petals.

*Proof (continued).* Reminder:  $p \ge 3$ ,  $\ell \ge 2$ ; true for smaller values of  $\ell$ ;  $\mathscr{D} \subseteq \mathscr{A}$  be a collection of pairwise disjoint sets of maximum possible size;  $D = \bigcup \mathscr{D}$ .

**Claim.** There exists some  $d \in D$  s.t. d belongs to more than  $(p-1)^{\ell-1}(\ell-1)!$  elements of  $\mathscr{A}$ .

Let  $d \in D$  be as in the Claim, and set  $\mathscr{A}' := \{A \setminus \{d\} \mid A \in \mathscr{A}, d \in A\}$ . Then  $|\mathscr{A}'| > (p-1)^{\ell-1}(\ell-1)!$ , and  $|A| \leq \ell - 1 \quad \forall A \in \mathscr{A}'$ . So, by the ind. hyp.,  $\exists$  sunflower  $\mathscr{S}' \subseteq \mathscr{A}'$  with p petals. Now, set  $\mathscr{S} := \{A \cup \{d\} \mid A \in \mathscr{S}'\}$ ; then  $\mathscr{S} \subseteq \mathscr{A}$  is a sunflower with p petals, and we are done.